

Dose-response effects of indomethacin and PGE₂ on electromechanical activity of in vivo rabbit ileum

K. L. KOCH, A. DWYER, AND G. H. JEFFRIES

Gastroenterology Division, Department of Medicine, The Milton S. Hershey Medical Center of Pennsylvania State University, Hershey, Pennsylvania 17033

KOCH, K. L., A. DWYER, AND G. H. JEFFRIES. *Dose-response effects of indomethacin and PGE₂ on electromechanical activity of in vivo rabbit ileum.* Am. J. Physiol. 250 (Gastrointest. Liver Physiol. 13): G135-G139, 1986.—We determined dose-response characteristics of indomethacin and prostaglandin E₂ on the myoelectric activity in ileum of anesthetized New Zealand White rabbits. Monopolar electrodes and an intraluminal saline-filled catheter were used to simultaneously record electrical and mechanical activity. Thirty minutes after injection of 3.0, 5.0, and 10.0 mg/kg indomethacin, the percentage of slow waves with action potentials increased significantly from 10% to over 80%; at 60 min action potential activity decreased but remained dose dependent and significantly greater than controls. Action potential activity correlated with phasic increases in intraluminal pressure. Low-dose indomethacin (1.5 mg/kg) did not significantly alter action potential activity. Action potential activity induced by indomethacin (5 mg/kg) decreased dose dependently after infusion of prostaglandin E₂ (PGE₂, 1–28 μg/kg). In summary, dose-dependent action potential activity was induced by indomethacin and reversed by PGE₂. Endogenous inhibitory prostaglandins (PGE₂ or others) appear to modulate activity of specific excitatory neuromuscular circuits in in vivo ileum.

intestinal motility; anti-inflammatory drugs; prostaglandins

INDOMETHACIN is a potent inhibitor of prostaglandin synthesis (20). A variety of prostaglandins are produced in gut tissues, including the microsomes of smooth muscle cells (19). The effects of indomethacin on intestinal smooth muscles have been extensively studied in vitro: indomethacin inhibits longitudinal muscle contraction in a variety of conditions (2, 5, 8, 10, 16); conversely, indomethacin stimulates contraction of circular muscle (6, 18, 21). The effects of indomethacin on in vitro ileal circular muscle strips are due to inhibition of prostaglandin production, particularly prostaglandin E₂ (PGE₂) and prostacyclin (PGI₂) (18). On the basis of in vitro studies, it has been proposed that endogenous prostaglandin synthesis has a physiological role in propulsive and nonpropulsive movements of the intestine (24) and in the genesis of intestinal tone (9).

Few studies, however, have examined the effects of indomethacin on in vivo myoelectric activity of the small intestine. Although changes in intraluminal pressure and action potential activity have been individually described after administration of indomethacin (13, 15, 20), a systematic evaluation of the effects of indomethacin on in

vivo electromechanical activity of the intestine has not been reported. The present study was undertaken to investigate the effects of indomethacin on in vivo ileum by simultaneously recording myoelectric and mechanical activity from rabbit ileum. The dose-response effects of PGE₂ on the myoelectric activity induced by indomethacin were also examined.

MATERIALS AND METHODS

Surgical preparation. Male New Zealand White rabbits weighing 2.0–2.9 kg were used in all studies. Water but not food was allowed ad libitum for 18 h prior to the experiments. Pentobarbital sodium (30 mg/kg) was administered through an ear vein. A tracheostomy was performed and a catheter was placed into the internal jugular vein for administration of indomethacin or additional pentobarbital. Body temperature was maintained at 38°C throughout each experiment by a heating blanket. Through a midline abdominal incision, the ileum was identified. Three monopolar silver-silver chloride electrodes were placed perpendicular to the longitudinal axis of the ileum at 2.5-cm intervals and sewn to the serosa and outermost layers of the longitudinal muscle. The electrodes were connected to a rectilinear recorder (Beckman, R612, Fullerton, CA) through 8353A couplers. A reference electrode was placed in the hindlimb of the rabbit. All recordings were made with settings yielding a sensitivity of 0.05 mV/mm with a high-frequency cutoff of 30 Hz and low-frequency filter of 0.16 Hz. To record intraluminal pressure, a saline-filled, open-tipped polyurethane catheter (PEG-190) was inserted into the lumen of the ileum through a purse-string suture 5 cm orad to the proximal electrode, positioned beneath a serosal electrode, and connected to a Beckman pressure transducer (type 4-327-C).

Experimental design. Control rabbits were prepared as described above, and after a 90-min equilibration period a 2-ml bolus of saline was administered intravenously. Myoelectric activity was recorded for the subsequent 5 h. After the equilibration period, a single experimental animal received a single dose of indomethacin (either 1.5, 3.0, 5.0, or 10.0 mg/kg).

Indomethacin (Merck Sharp & Dohme, West Point, PA) and sodium bicarbonate (3:1 ratio by weight) were dissolved in saline and injected intravenously in a 2-ml bolus. PGE₂ (Sigma Chemical, St. Louis, MO) was dissolved in ethanol, evaporated under nitrogen, and dis-

solved in physiological phosphate buffer. Thirty to sixty minutes after injection of indomethacin (5 mg/kg), PGE₂ was administered intravenously over 30 s at doses ranging from 1 to 28 µg/kg.

Myoelectric data. Myoelectric recordings were scored by hand for slow wave frequency per minute and the percentage of slow waves with action potentials. Myoelectric activity from electrodes yielding the clearest slow waves throughout the experiment were selected for analysis. Slow wave frequency and the percentage of slow waves with action potentials were assessed during the 10-min period immediately preceding the injection of indomethacin and the 20- to 30-min and 50- to 60-min periods after injection. The same myoelectric characteristics were determined during 10-min periods before and after injection of PGE₂. Data were recorded as means ± SE.

Statistical analysis. Slow wave frequencies and percentages of slow waves with action potentials from control and indomethacin experiments were compared by use of unpaired *t* tests. Paired *t* tests were used to determine the significance of myoelectric effects recorded after PGE₂. A value of *P* < 0.05 was considered significant.

RESULTS

Myoelectric activity. Action potentials occurred on 0–7% of the slow waves recorded from control rabbits, and simultaneously recorded intraluminal pressures were quiescent. This electromechanical pattern was recorded for at least 5 h in the control rabbits.

Typical electromechanical changes observed after injection of indomethacin (5 mg/kg) are shown in Fig. 1. Action potential activity commenced after a lag time of 13–18 min (mean 14.8 ± 1.0 min, *n* = 5) after indomethacin. The onset of action potential activity was progressive, and maximum action potential activity occurred between 30 and 60 min after injection of indomethacin. Action potential activity was recorded at each electrode site and appeared on discrete regions of the slow waves. Phasic increases in intraluminal pressure correlated in time with the onset of action potentials on individual slow waves. The amplitude of intraluminal pressures ranged from 10 to 15 mmHg. Visual inspection of the ileum revealed intermittent contractions during intense action potential activity induced by indomethacin.

The dose-response characteristics of indomethacin on action potential activity are shown in Fig. 2. Action potential activity for control and experimental rabbits was equivalent at *time 0*. Indomethacin, 3.0, 5.0, and 10.0 mg/kg, significantly increased the percentage of slow waves with action potentials when compared with controls at 30 min. In contrast, 1.5 mg/kg indomethacin had an insignificant effect on action potential activity at 30 min. The maximum increase in the percentage of slow waves with action potential activity observed in these experiments (81 ± 9%) occurred 30 min after 5 mg/kg indomethacin; 10 mg/kg indomethacin did not further increase action potential activity (62.2 ± 8 at 30 min). At 60 min, action potential activity decreased but re-




FIG. 1. Electromechanical recording from rabbit ileum 60 min after injection of indomethacin (5 mg/kg iv). Electrodes (E₁–E₃) are 2.5 cm apart, positioned perpendicular to longitudinal axis of ileum and located proximal (E₁) to distal (E₃) on ileal serosa. P₂ indicates intraluminal pressure recorded beneath E₂. Peak pressures point to bottom of tracing. Slow wave frequency is 16 cycles/min. Action potential activity (AP), particularly well seen at E₂, occurs on discrete portions of slow waves recorded at each electrode site. Onset of action potential activity on individual slow waves at E₂ corresponds with phasic increases in intraluminal pressure at P₂.




FIG. 2. Dose-response effect of indomethacin on percentage of slow waves (±SE) with action potentials before indomethacin (*time 0*) and 30 and 60 min after injection. Indomethacin significantly increases action potential activity. *†Significant differences between saline and indomethacin values. See text for description. Number of rabbits receiving each dose is shown on Table 1.

mained significantly greater than base-line activity after 3.0, 5.0, and 10 mg/kg indomethacin.

Base-line slow wave frequencies were not significantly altered 30 and 60 min after injection of either saline or

the various doses of indomethacin (Table 1). However, when slow wave frequencies from control and indomethacin-treated rabbits were compared (unpaired *t* tests), significantly lower frequencies were found 30 min after 1.5 and 3.0 mg/kg indomethacin and 60 min after 3.0 and 5.0 mg/kg indomethacin. In contrast, slow wave frequencies after 10 mg/kg indomethacin did not differ from control. The overall effect of indomethacin on slow wave frequency was small, and a dose-response effect was not apparent.

Effect of PGE₂ on indomethacin-induced electromechanical activity. In other experiments, PGE₂ (1–28 μg/kg) was injected intravenously 30–60 min after administration of 5 mg/kg indomethacin at a time when action potential activity occurred on at least 60% of the slow waves. The effect of each dose of PGE₂ was evaluated during the 10-min periods before and after injection in a minimum of four rabbits. PGE₂ decreased the indomethacin-induced action potential activity in a dose-dependent manner as shown in Fig. 3. PGE₂, 1 μg/kg, had no effect on action potential activity; 4 and 6 μg/kg PGE₂ reduced the percentage of slow waves with action potentials but not significantly. After 8 μg/kg PGE₂, the action

TABLE 1. Effect of indomethacin on slow wave frequency in *in vivo* rabbit ileum

	n	Slow Wave Frequency		
		0 min	30 min	60 min
Control	5	17.9±0.6	18.2±0.5	18.4±0.6
Indomethacin, mg/kg				
1.5	4	15.5±0.6	15.4±0.4*	15.6±0.5
3.0	4	15.9±0.7	15.4±0.6*	16.0±0.6†
5.0	5	17.3±0.5	16.5±0.6	15.9±0.3*
10.0	5	15.8±0.5	15.9±0.9	16.1±1.0

Values are means ± SE in cycles/min; n, no. of measurements. **P* < 0.01 Indo vs. control. †*P* < 0.02 Indo vs. control.




FIG. 3. Dose-response effects of PGE₂ on percentage of slow waves with action potentials (±SE) induced by indomethacin. PGE₂ was administered 30–60 min after injection of indomethacin (5 mg/kg) when action potentials were occurring on 60% or more of slow waves (Before PGE₂). Increasing doses of PGE₂ abolish indomethacin-induced action potentials (After PGE₂). Number of rabbits receiving each dose is shown on Table 2. *Significant differences between effects of 1 μg/kg PGE₂ and other PGE₂ doses.

TABLE 2. Effect of PGE₂ on slow wave frequency in rabbits pretreated with 5 mg/kg indomethacin

PGE ₂ , μg/kg	n	Slow Wave Frequencies		
		Before PGE ₂	After PGE ₂	<i>P</i>
1	4	16.1±0.2	15.9±0.2	NS
4	5	16.3±0.2	16.1±0.4	NS
6	5	17.7±0.4	18.8±0.6	NS
8	9	16.7±0.4	19.1±0.6	<0.01
18	8	17.0±0.3	19.7±0.5	<0.01
28	7	17.5±0.7	19.3±0.3	<0.01

Values are means ± SE in cycles/min.

potential activity was significantly reduced to 13 ± 7%; after 18 and 28 μg/kg action potential activity was virtually abolished (5 ± 5 and 4 ± 2%, respectively). Action potential activity and phasic increases in intraluminal pressure were temporarily abolished by 18 and 28 μg/kg PGE₂; the duration of the inhibitory PGE₂ effect was 10–12 min.

Slow wave frequencies 10 min before and after the various doses of PGE₂ are listed in Table 2. PGE₂ at 1, 4, and 6 μg/kg did not affect slow wave frequency, whereas 8, 18, and 28 μg/kg significantly increased the slow wave frequency in the indomethacin-treated rabbits. In other experiments (*n* = 6), a dose 18 or 28 μg/kg PGE₂ was administered after the equilibration period but before indomethacin. Neither slow wave frequency nor action potential activity was altered by the two PGE₂ doses. Combined data from these experiments are listed below: slow wave frequency before and after PGE₂ was 19.0 ± 0.4 and 19.1 ± 0.3, respectively, and the percentage of slow waves with action potential activity was 5.0 ± 2.0 and 4.4 ± 1.0, respectively.

Histological examination. Histological examination of ileum from control and indomethacin-treated rabbits revealed no mucosal or muscle wall abnormalities by routine light microscopy.

DISCUSSION

Endogenous prostaglandins may have a physiological role in modulating intestinal motility (5, 9, 24). By altering biosynthesis of prostaglandins with indomethacin and measuring subsequent electromechanical activity, the effect of ongoing endogenous production of prostaglandins on intestinal motility may be assessed (18, 20). Although indomethacin is a well-known and potent inhibitor of prostaglandin synthesis (22), a detailed examination of the effects of indomethacin on *in vivo* electromechanical activity of the intestine has not been reported. The present *in vivo* studies show that indomethacin induces dose-dependent action potential activity that is associated with phasic increases in intraluminal pressure. The lag time to onset of action potential activity after indomethacin is consistent with lag times reported for inhibition of cyclooxygenase and depletion of prostaglandins in ileal and antral smooth muscle (18, 20, 21). Although tissue prostaglandin levels were not measured in these experiments, Wilson et al. (25) previously showed that PGE was significantly reduced in rabbit intestine after 5 mg/kg indomethacin, the most active dose used in the present studies.

The *in vivo* electromechanical activity observed after indomethacin is characterized by rhythmic bursts of action potentials on slow waves and phasic increases in intraluminal pressure, all of which indicate contractile activity of the circular muscle layer (1). Similar action potential activity after indomethacin has been noted in the jejunum of fasted conscious dogs (13). Thus, anesthesia and laparotomy *per se* do not appear relevant to the electromechanical activity noted after indomethacin. Indomethacin also stimulates contractions of *in vitro* circular muscle from human and canine ileum (6, 18, 19). The increase in *in vitro* muscle contractility is associated with decreased tissue levels of PGE₂ and PGI₂, changes which occurred after a lag phase of 20 min duration (18). Similar lag periods preceded the onset of indomethacin-induced electromechanical activity in the present *in vivo* studies. Taken together, these data suggest that the dose-dependent *in vivo* action potential activity observed after indomethacin was due to depletion of inhibitory endogenous prostaglandins that modulate circular muscle contractility.

Administration of PGE₂ resulted in dose-dependent reversal of the *in vivo* action potential activity induced by indomethacin. It is well documented that PGE₂ inhibits contraction of circular muscle *in vitro* (3, 4, 21). Other studies have shown that PGE₂ decreases intraluminal pressure, an effect interpreted as inhibition of circular muscle contraction (3, 7). The inhibition of *in vivo* action potential activity (i.e., circular muscle contraction) by PGE₂ is consistent with these previous studies. Furthermore, *in vitro* circular muscle contractions induced by indomethacin are also inhibited by PGE₂ and PGI₂ (18). It has been previously proposed that PGE₂ and PGI₂ inhibit potential excitatory neuromuscular activity in the circular muscle (13, 18, 20). Although PGI₂ was not studied, the effects of indomethacin and PGE₂ reported here strongly support the concept that inhibitory endogenous prostaglandins such as PGE₂ modulate intestinal motility. Specifically, the present dose-response experiments suggest that inhibitory prostaglandins regulate the activity of excitatory neuromuscular circuits, which mediate phasic contractions of *in vivo* ileal circular muscle.

Aspirin (100 mg/kg *iv*) produced electromechanical activity in rabbit ileum which was similar to indomethacin-induced responses (12). These findings indicate that indomethacin-induced action potential activity is the result of inhibition of prostaglandin synthesis rather than a nonspecific effect of the drug itself. Thus, by inhibiting endogenous prostaglandins indomethacin may unmask excitatory neuromuscular pathway(s) that are ordinarily suppressed by the ongoing synthesis of prostaglandin products such as PGE₂ or PGI₂ (13, 18, 20). The dose-dependent reversal of indomethacin-induced action potential activity by PGE₂ supports this notion. Furthermore, previous studies showed that indomethacin-induced action potentials were significantly decreased by scopolamine but were unaffected by hexamethonium, phentolamine, propranolol, and bilateral cervical vagotomy (11). Thus, indomethacin appears to unmask an enteric cholinergic circuit in rabbit ileum by

inhibiting local synthesis of prostaglandins. The dose-dependent effects of indomethacin on *in vivo* action potential activity support the concept that a distinct enteric neuromuscular circuit(s) or receptor(s) mediates the electromechanical response observed after indomethacin.

The effect of indomethacin and PGE₂ on *in vivo* slow wave frequency deserves comment. Slow waves probably originate in the longitudinal muscle layer of the intestine, generated by metabolic processes within the muscle that affect membrane depolarization and repolarization (23). In the present studies some doses of indomethacin caused a slight decrease in slow wave frequency, but a dose-response effect was not seen. Thus, indomethacin would appear to have little effect on the longitudinal muscle. However, if indomethacin induced a deficiency of prostaglandins and altered cellular processes that affected slow waves, then administration of prostaglandin should reverse the effect. As shown in Table 2, infusions of PGE₂ resulted in significant dose-dependent increases in slow wave frequency, an effect observed only in indomethacin-treated rabbits. Increased slow wave frequency was also noted in *ex vivo* canine stomach after flash or continuous infusions of PGE₂, a finding attributed to selective action on the longitudinal muscle (14). Thus, prostaglandins may also have a role in the generation of *in vivo* slow waves through as-yet-undefined metabolic effects in the longitudinal muscle.

The authors acknowledge Evelyn Lewis and Joyce Zeager for their excellent secretarial assistance.

The investigators adhered to the "Guide for the Care and Use of Laboratory Animals" as prepared by the Committee on the Guide for Laboratory Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences, National Research Council.

Address for reprint requests: K. L. Koch, Gastroenterology Division, Milton S. Hershey Medical Center, Pennsylvania State University, PO Box 850, Hershey, PA 17033.

Received 20 February 1985; accepted in final form 27 August 1985.

REFERENCES

- BASS, P., AND J. N. WILEY. Electrical and extraluminal contractile force activity of the duodenum of the dog. *Am. J. Dig. Dis.* 10: 183-200, 1965.
- BENNETT, A., K. G. ELEY, AND H. L. STOCKLEY. Modulation by prostaglandins of contractions in guinea-pig ileum. *Prostaglandins* 9: 377-384, 1975.
- BENNETT, A., K. G. ELEY, AND G. B. SHOLES. Effect of prostaglandins E₁ and E₂ on intestinal motility in the guinea-pig and rat. *Br. J. Pharmacol.* 34: 639-647, 1968.
- BENNETT, A., AND B. FLESHLER. Prostaglandins and the gastrointestinal tract. *Gastroenterology* 59: 790-800, 1970.
- BOTTING, J. H., AND R. SALZMANN. The effect of indomethacin on the release of prostaglandin E and acetylcholine from guinea-pig isolated ileum at rest and during field stimulation. *Br. J. Pharmacol.* 50: 119-124, 1974.
- BURLEIGH, D. E. The effects of indomethacin on the tone and spontaneous activity of the human small intestine *in vitro*. *Arch. Int. Pharmacodyn. Ther.* 225: 240-245, 1977.
- DAJANI, E. Z., E. A. W. ROGE, AND R. E. BERTERMANN. Effects of E prostaglandins, diphenoxylate and morphine on intestinal motility *in vivo*. *Eur. J. Pharmacol.* 34: 105-113, 1975.
- EHRENPREIS, S., AND J. GREENBERG. Prostaglandins reverse inhibition of electrically-induced contractions of guinea pig ileum by morphine, indomethacin and acetylsalicylic acid. *Nature Lond.* 245: 280-282, 1973.

9. FERREIRA, S. H., A. HERMAN, AND J. R. VANE. Prostaglandin generation maintains the smooth muscle tone of the rabbit isolated jejunum (Abstract). *Br. J. Pharmacol.* 44: 328P-330P, 1972.
10. KADLEC, O., K. MASEK, AND I. SEFERNA. A modulating role of prostaglandins in contractions of the guinea-pig ileum. *Br. J. Pharmacol.* 51: 565-570, 1974.
11. KOCH, K. L., AND A. DWYER. Indomethacin-induced spiking activity in rabbit ileum: in vivo interactions of prostaglandins and cholinergic neurotransmission. In: *Proceedings of the 9th International Symposium on Gastrointestinal Motility*, edited by C. Roman. Lancaster, UK: 1983, p. 357-363.
12. KOCH, K. L., A. DWYER, AND G. H. JEFFRIES. Electromechanical effects of acetylsalicylic acid and sodium salicylate on *in vivo* rabbit ileum (Abstract). *Clin. Res.* 31: 284A, 1983.
13. KONTUREK, S. J., P. THOR, W. PAWLIK, P. GUSTAW, AND A. DEMBINSKI. The role of prostaglandins in the myoelectric, motor and metabolic activity of the small intestine in the dog. In: *Motility of the Digestive Tract*, edited by M. Wienbeck. New York: Raven, 1982, p. 437-444.
14. KOWALEWSKI, K., AND A. KOLODEY. Effect of prostaglandin-E₂ on myoelectric and mechanical activity of totally isolated, *ex vivo*-perfused, canine stomach. *Pharmacology* 13: 325-339, 1975.
15. MATHIAS, J. R., G. M. CARLSON, G. BERTIGER, J. L. MARTIN, AND S. COHEN. Migrating action potential complex of cholera: a possible prostaglandin-induced response. *Am. J. Physiol.* 232 (*Endocrinol. Metab. Gastrointest. Physiol.* 1): E529-E534, 1977.
16. NORTHOVER, B. J. Mechanism of the inhibitory action of indomethacin on smooth muscle. *Br. J. Pharmacol.* 41: 540-551, 1971.
17. OUYANG, A., C. J. CLAIN, W. J. SNAPE, AND S. COHEN. Characterization of opiate-mediated responses of the feline ileum and ileocecal sphincter. *J. Clin. Invest.* 69: 507-515, 1983.
18. SANDERS, K. M. Evidence that prostaglandins are local regulatory agents in canine ileal circular muscle. *Am. J. Physiol.* 246 (*Gastrointest. Liver Physiol.* 9): G361-G371, 1984.
19. SANDERS, K. M., AND T. E. NORTHRUP. Prostaglandin synthesis by microsomes of the circular and longitudinal gastrointestinal muscles. *Am. J. Physiol.* 244 (*Gastrointest. Liver Physiol.* 7): G442-G448, 1983.
20. SANDERS, K. M., AND G. ROSS. Effects of endogenous prostaglandin E on intestinal motility. *Am. J. Physiol.* 234 (*Endocrinol. Metab. Gastrointest. Physiol.* 3): E204-E208, 1978.
21. SANDERS, K. M., AND J. H. SZURSZEWSKI. Does endogenous prostaglandin affect gastric antral motility? *Am. J. Physiol.* 241 (*Gastrointest. Liver Physiol.* 4): G191-G195, 1981.
22. VANE, J. R. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nature Lond.* 231: 232-235, 1971.
23. WEISBRODT, N. W. Motility of the small intestine. In: *Physiology of the Gastrointestinal Tract*, edited by L. R. Johnson. New York: Raven, 1981, p. 411-443.
24. WILSON, D. E. Prostaglandins. Their actions on the gastrointestinal tract. *Arch. Intern. Med.* 133: 112-118, 1974.
25. WILSON, D. E., S. EL-HINDI, P. TAO, AND L. POPPE. Effects of indomethacin on intestinal secretion, prostaglandin E and cyclic AMP: evidence against a role for prostaglandin E and cyclic AMP: evidence against a role for prostaglandins in cholera toxin-induced secretion. *Prostaglandins* 10: 581-587, 1975.

